



ORIGINAL ARTICLE

# Mirror therapy in chronic stroke survivors with severely impaired upper limb function: a randomized controlled trial

Carolina COLOMER<sup>1\*</sup>, Enrique NOÉ<sup>1</sup>, Roberto LLORENS<sup>1,2</sup>

<sup>1</sup>Servicio de Neurorrehabilitación y Daño Cerebral de los Hospitales NISA, Fundación Hospitales NISA, Valencia, Spain; <sup>2</sup>Instituto Interuniversitario de Investigación en Bioingeniería y Tecnología Orientada al Ser Humano, Universitat Politècnica de València, Valencia, Spain

\*Corresponding author: Carolina Colomer, Hospital Nisa Valencia al Mar. Río Tajo 1, 46011 Valencia, Spain. E-mail: [carol@neurorhb.com](mailto:carol@neurorhb.com)

## ABSTRACT

**BACKGROUND:** Mirror therapy (MT) has been proposed to improve the motor function of chronic individuals with stroke with mild to moderate impairment. With regards to severe upper limb paresis, MT has shown to provide limited motor improvement in the acute or sub-acute phase. However, no previous research has described the effects of MT in chronic individuals with stroke with severely impaired upper limb function.

**AIM:** The aim of this study was to determine the effectiveness of MT on chronic stroke survivors with severe upper-limb impairment in comparison with passive mobilization.

**DESIGN:** A randomized controlled trial.

**SETTING:** Rehabilitative outpatient unit.

**POPULATION:** A total of 31 chronic subjects poststroke with severely impaired upper limb function were randomly assigned to either an experimental group (N.=15), or a control group (N.=16).

**METHODS:** Twenty-four intervention sessions were performed for both groups. Each session included 45-minute period of MT (experimental group) or passive mobilization (control group), administered three days a week. Participants were assessed before and after the intervention with the Wolf Motor Function Test, the Fugl-Meyer Assessment, and the Nottingham Sensory Assessment.

**RESULTS:** Improvement in motor function was observed in both groups on the time (P=0.002) and ability (P=0.001) subscales of the Wolf Motor Function Test. No differences were detected in kinesthesia or stereognosis. However, the experimental group showed a significant improvement in tactile sensation that was mainly observed as an increased sensitivity to light touches.

**CONCLUSIONS:** In comparison with passive mobilization, MT in chronic stroke survivors with severely impaired upper-limb function may provide a limited but positive effect on light touch sensitivity while providing similar motor improvement.

**CLINICAL REHABILITATION IMPACT:** MT is a therapeutic approach that can be used in the rehabilitation of severely impaired upper limb in chronic stroke survivors, specifically to address light touch sensitivity deficits.

*(Cite this article as: Colomer C, Noé E, Llorens R. Mirror therapy in chronic stroke survivors with severely impaired upper limb function: a randomized controlled trial. Eur J Phys Rehabil Med 2016;52:271-8)*

**Key words:** Stroke - Upper Extremity - Paresis - Brain injury, chronic.

Functional impairment of the upper limb is reported in approximately 85% of stroke survivors<sup>1</sup> and affects participation in daily living activities and quality of life.<sup>2</sup> Six months after onset, 30-60% of individuals do not regain functional use, and only 5-20% will achieve full recovery of arm function.<sup>3</sup> Rehabilitation of severe arm paresis in chronic stroke survivors is, therefore, especially challenging.<sup>4</sup> Useful reorganization of cortical

areas involved in arm function occurs in response to active exercise and to motor and attentional inclusion of the affected arm in task oriented movements.<sup>4,5</sup> Because severe paresis impedes active training of the hand, traditionally, rehabilitation strategies in these subjects have almost exclusively aimed to compensate for the deficit by training the opposite limb in daily tasks.<sup>4,6</sup> As synaptic connectivity is highly use dependent, this ab-

sense of stimulation on the chronic paretic arm results in reduced sensorimotor representation in the available neural circuits over time<sup>7</sup> and consequently diminishes the possibilities for sensorimotor clinical progress.<sup>6</sup> Accordingly, a reduction in sensorimotor abilities might partly be an effect of non-use of the affected limb.<sup>7</sup> In fact, lack of movement has been considered to be a form of “learned paralysis”.<sup>8,9</sup>

Mirror therapy (MT) could address these issues because it does not require active movements of the affected arm. A mirror is placed along the midsagittal plane between the two limbs, and the subject is encouraged to move the less affected limb while watching its reflection in the mirror, thus providing the visual illusion that the affected limb mirrors the movement of the other limb. Neuroimaging techniques have revealed the capacity of MT to elicit cortical activation in the hemisphere contralateral to the mirrored arm (ipsilateral to the moving arm), even in absence of movement,<sup>10</sup> and have found evidence of interesting and varied cortical changes induced by MT.<sup>11-13</sup> These results have motivated the application of MT to sensorimotor rehabilitation after stroke. Previous research has shown that MT can improve the motor function of chronic individuals with stroke with mild to moderate impairment.<sup>14, 15</sup> Moreover, interventions involving stroke survivors with severe upper limb paresis have been shown to provide limited motor improvement in the acute<sup>10, 16</sup> or sub-acute phase.<sup>16</sup> However, no previous research has described the effects of MT in chronic individuals with stroke with severely impaired upper limb function. We hypothesize that MT can be beneficial for sensorimotor function of severely affected upper limbs of chronic stroke survivors.

The objective of the present study was to determine the effectiveness of MT in chronic stroke survivors with severe upper-limb impairment in comparison to a passive mobilization intervention.

## Materials and methods

### Participants

Participants were recruited from the chronic stroke outpatient management program that was run by a specialized neurorehabilitation center. The inclusion criteria for the current study were: 1) chronicity > six months; 2) severe paresis of the upper limb, as defined by the

Brunnstrom Approach<sup>17</sup> as stages I or II) and by the Upper Extremity subscale of the Fugl-Meyer Assessment (FMA)<sup>18</sup> as scores below 19; 3) sensory impairment in the affected upper limb, as assessed by clinical examination; 4) ability to maintain a sitting position for at least 60 minutes; 5) a fairly good cognitive condition, as defined by scores on the Mini-Mental State Examination<sup>19</sup> above 23. Participants were excluded if they had: 1) impaired comprehension that hindered sufficient understanding of the instructions, as defined by Mississippi Aphasia Screening Test<sup>20</sup> scores below 45; 2) severe visual impairments; 3) upper limb pain that limited participation in the rehabilitation protocol; 4) spatial neglect; 5) self-awareness disorders; and 6) emotional circumstances that impeded adequate collaboration. Sample size was calculated for a two-sample *t*-test assuming 75% power, a common standard deviation of 50, a mean difference between groups on the Wolf Motor Function Test (WMFT) of 50 seconds, and a loss rate of 10%.

The study was approved by the Institutional Review Board of the specialized neurorehabilitation center. Written consent was obtained from all of the subjects who satisfied the inclusion criteria and accepted the offer to participate in the study. Participants were randomly assigned to an experimental or a control group. The allocation sequence was concealed from an independent researcher. A sealed envelope identifying the group of each participant was given to the therapists to inform them of the allocation. Randomization was computer-generated using a basic random number generator in a ratio of 1:1.

### Intervention protocol

All of the participants were undergoing a long-term care physical therapy program consisting of five one-hour sessions a week that focused on balance and gait training. During this clinical trial, participants underwent 24 sessions, each 45-minutes long, and administered three days a week in addition to the physical therapy. Subjects belonging to the experimental group underwent a MT program and those belonging to the control group received passive mobilization of the affected upper limb. Treatment intensity was dose-matched for both groups, allowing 60-second breaks every five minutes. All of the participants trained in a dedicated area of the physical therapy unit free of dis-

tractors. For the MT, a triangular prism-shaped device with a mirror on one side was fixed on a conventional table. Participants sat down close to the table in chairs with a back and without armrests. Participants hid the affected arm inside the device and the mirror was oriented so that they were able to see the reflection of their other arm. Participants were encouraged to observe the mirror while executing different movements with their less affected limb. Exercises were indicated verbally and consisted of a series of flexo-extension and pronosupination movements of the shoulder and forearm and gross and fine movements of the wrist, hand, and fingers, with and without objects (balls, cups, and other). These activities included transitive movements, gross motor tasks, and intransitive movements.<sup>14</sup> Participants in the control group received passive mobilization of the paretic upper limb. Passive range of motion exercises were provided in those segments where no active movement was detected to meticulously reproduce a range of articular movements and muscle and soft tissue elongation. In case of residual active movement capability, participants were encouraged to perform the movements with the assistance of the therapists.

### Assessment

All of the participants were assessed by a physical therapist, blind to the treatment, the day before the intervention and the day after the intervention. The primary outcome measure was the WMFT. Both time and ability subscales were considered. Secondary outcome measures assessed the upper limb motor function using the FMA, and the sensory impairment of the hemiparetic upper extremity using the tactile, kinesthetic, and stereognosis subscales of the Nottingham Sensory Assessment (NSA).<sup>21</sup> The tactile subscale assessed light touch, pressure, pinprick, temperature, tactile localization, and bilateral simultaneous touch. Scores in the NSA ranged from 0 to 48 in the tactile subscale, from 0 to 12 in the kinesthetic subscale, and from 0 to 22 in the stereognosis subscale. Higher scores represent more preserved sensory sensation.

### Statistical analysis

Demographical and clinical comparisons between the control and the experimental groups were performed

with independent sample *t*-tests and  $\chi^2$  or Fisher exact tests, as appropriate. Repeated measures analyses of variance (ANOVA) with time as the within-subjects factor and treatment option (control versus experimental) as the between-subjects factor were performed for the FMA and the subscales of the WFMT and NSA. The main effects were evaluated for time, treatment option, and the time-treatment option interaction. ANOVA findings that violated the sphericity assumption were accommodated by Greenhouse and Geisser's conservative degrees of freedom adjustment. For each repeated-measures ANOVA, we present the partial eta squared ( $\eta^2_p$ ) as a measure of effect size; values may range between 0 and 1, with higher values representing higher proportions of variance explained by the independent variable.

The  $\alpha$  level was set at 0.05 for all analyses (two-sided). All analyses were computed with SPSS for Mac, version 15 (IBM, Armonk, NY, USA). Investigators performing the data analysis were blinded.

## Results

A total pool of 97 outpatients attended the long-term care program during the recruitment. Of those, 34 subjects (35.1%) met inclusion criteria. None of them refused to participate in the study, and consequently all of them were randomized. Each group consisted of 17 participants. Two participants of the experimental group were discharged, and one participant of the control group suffered a cardiac arrest. Consequently, these participants discontinued the program and were excluded from the study. Their data were, therefore, not included for analysis. Finally, data from 31 participants, 16 in the control group and 15 in the experimental group, were included (Figure 1). The final sample consisted of 26 males and 5 females, with a mean age of  $53.6 \pm 8.4$  years, and a mean chronicity of  $551.1 \pm 377.5$  days. A total of 23 participants presented an ischemic stroke, and eight participants presented a hemorrhagic stroke (Table I). No significant differences in demographical (gender and age) or clinical (etiology, hemiparetic side, chronicity, Brunnstrom, and FMA) data at inclusion were detected between the groups. All of the participants were right-handed.

With regards to the primary outcome measure, scores in the time and ability subscales of the WMFT significantly improved after the treatment period in each




Figure 1.—CONSORT flow diagram. Progress through the phases of the parallel randomized trial of both groups.

group, but this improvement failed to reach a significant difference between groups (Table II). The percentage of change was 5.8% in the control group and 4.7% in the experimental group for the time subscale, and 15.6% in the control group and 18.9% in the experimental group for the ability subscale.

Regarding to secondary outcomes, no differences were found in motor performance after the treatment, as measured by the FMA, neither within any group nor be-

tween the two groups (Table II). Regarding sensory assessment, the kinesthetic and the stereognosis subscales of the NSA showed no significant difference (Table II). However, the results in the tactile subscale revealed a statistically significant improvement after treatment for both groups. Interestingly, the improvement experienced by the experimental group was significantly higher than that experienced by the control group ( $P < 0.01$ ). A more in-depth analysis of the tactile subscales found

This document is protected by international copyright laws. No additional reproduction is authorized. It is permitted for personal use to download and save only one file and print only one copy of this Article. It is not permitted to make additional copies (either sporadically or systematically, either printed or electronic) of the Article for any purpose. It is not permitted to distribute the electronic copy of the article through online internet and/or intranet file sharing systems, electronic mailing or any other means which may allow access to the Article. The use of all or any part of the Article for any Commercial Use is not permitted. The production of derivative works from the Article is not permitted. It is not permitted to remove, cover, overlay, obscure, block, or change any copyright notices or terms of use which the Publisher may post on the Article. It is not permitted to frame or use framing techniques to enclose any trademark, logo or other proprietary information of the Publisher.

TABLE I.—*Characteristics of the participants.*

Characteristic (N.=15)	Control group (N.=16)	Experimental group	Significance
Gender (n, %)			NS (P=0.532)
Male	13 (81.2%)	13 (86.7%)	
Female	3 (18.8%)	2 (13.3%)	
Age (years)	53.3±10.5	53.8±5.5	NS (P=0.891)
Etiology (n, %)			NS (P=0.352)
Ischemic stroke	13 (81.2%)	10 (66.7%)	
Hemorrhagic stroke	3 (18.8%)	5 (33.3%)	
Hemiparesis (n, %)			NS (P=0.173)
Left	14 (87.5%)	10 (66.7%)	
Right	2 (12.5%)	5 (33.3%)	
Chronicity (days)	520.0±262.5	584.2±478.7	NS (P=0.601)
Brunnstrom (n, %)			NS
Stage I	0	0	
Stage II	16	15	
Flugl-Meyer Assessment	9.0±4.4	8.5±4.7	NS (P=0.803)

Age and chronicity are defined in terms of mean and standard deviation. Etiology and gender are expressed as percentage of the total number of participants. NS: non-significant.

TABLE II.—*Clinical data.*

	Initial assessment	Final assessment	Significance (P, effect size)
WMF-time			T** (P=0.002, $\eta^2_p=0.29$ )
Control	1492.7±65.1	1405.8±70.8	
Experimental	1615.2±67.2	1539.8±72.8	
WMF-ability			T** (P=0.001, $\eta^2_p=0.31$ )
Control	10.9±1.7	12.6±1.8	
Experimental	8.7±1.7	10.1±1.8	
FMA			NS
Control	9.0±1.1	9.5±1.1	
Experimental	8.5±1.2	8.6±1.1	
NSA-tactile			T** (P=0.001, $\eta^2_p=0.38$ ); GxT* (P=0.027, $\eta^2_p=0.16$ )
Control	23.9±4.5	25.1±4.3	
Experimental	17.8±4.7	21.9±4.4	
NSA-kinaesthetic			NS
Control	4.7±0.8	5.0±0.9	
Experimental	6.6±0.8	6.3±1.0	
NSA-stereognosis			NS
Control	5.2±1.8	5.3±1.8	
Experimental	4.2±1.8	4.4±1.8	

Results are given in terms of mean and standard deviation. T: time effect; GxT: group by time effect; \*P<0.05, \*\*P<0.01; NS: non-significant.

that scores in the light touch test promoted this result (Table III). No significant differences were detected in any subscale of the NSA between subjects with left or right hemiparesis, neither at the beginning nor at the end of the intervention.

## Discussion

This paper describes the effectiveness of a MT intervention in chronic stroke survivors with severe hemiparesis

in comparison with passive mobilization. Although the MT intervention provided similar results to mobilization interventions in motor performance, the experimental intervention could promote greater changes in the tactile sensation, specifically regarding light touch.

## Motor function

The improvement provided by MT for motor function was similar to the improvement provided by the

TABLE III.—Results in the tactile subscale of the Nottingham Sensory Assessment.

	Initial assessment	Final assessment	Significance (P, effect size)
Light touch			T** (P=0.000, $\eta^2_p=0.43$ ); GxT** (P=0.000, $\eta^2_p=0.36$ )
Control	4.2±0.8	4.3±0.7	
Experimental	3.5±0.8	4.3±0.7	
Pressure			NS
Control	4.4±0.8	4.5±0.7	
Experimental	4.2±0.8	4.3±0.8	
Pinprick			T** (P=0.001, $\eta^2_p=0.34$ )
Control	4.3±0.8	4.6±0.7	
Experimental	3.8±0.9	4.3±0.7	
Temperature			T* (P=0.023, $\eta^2_p=0.17$ )
Control	4.2±0.8	4.5±0.7	
Experimental	3.3±0.8	4.0±0.8	
Tactile localization			NS
Control	4.8±0.8	5.6±0.8	
Experimental	3.8±1.2	4.0±1.1	
Bilateral simultaneous touch			NS
Control	6.1±0.9	6.0±0.9	
Experimental	5.2±1.3	5.6±1.3	

Results are given in terms of mean and standard deviation. T: time effect; GxT: group by time effect; \*P<0.05, \*\*P<0.01; NS: non-significant.

mobilization intervention. Significant changes were detected in both groups by the WMFT, but not the FMA. Changes detected in the time subscale of the WMFT for both groups were lower than the Minimally Clinically Important Difference (MCID),<sup>22</sup> and more importantly, final timed scores, though lower than initial scores, were still very high, as expected for subjects with such severe motor impairment. Changes in the ability subscale in the control group were slightly inferior to the MCID, which was previously established as 17% for the acute phase,<sup>22</sup> but were higher than the MCID in the experimental group. The general nature of the FMA could explain the inability of this scale to detect changes as subtle as those reported in our study. The characteristics of our sample could have prevented greater improvement. First, all of the participants were chronic stroke survivors. Although chronicity is not believed to be an excluding factor, less recovery is expected as time since injury increases and endogenous recovery mechanisms diminish.<sup>23</sup> Second, all of the participants presented a severe upper limb motor impairment, evidenced by a baseline FMA score below 19. Initial severity of motor impairment has been considered the most important prognostic variable of upper limb recovery after stroke.<sup>24-26</sup> Specifically, an FMA above 18 points at four weeks after onset is an independent predictor of dexterity within six months.<sup>24</sup> More importantly, initial

FMA scores between 21 and 35 can predict improvement after a MT intervention.<sup>27</sup> Finally, somatosensory impairment, which was present in all of our participants as well, has also been found to be a predictor of the absence of limb dexterity one year after the injury,<sup>25</sup> and has been related to poorer functional ability.<sup>28, 29</sup> Previous research with similar interventions involving stroke survivors in the acute<sup>10</sup> or sub-acute phase<sup>16</sup> with severe upper limb paresis reported no significant differences between groups in the motor domain, except for individuals with plegic fingers.<sup>10</sup> The combination of MT with bilateral arm training in an acute population was reported to increase the effectiveness of the intervention.<sup>30</sup> In contrast, positive effects were found in motor performance and motor control after MT in chronic individuals with mild to moderate hemiparesis.<sup>14, 15</sup> All of these results suggest that the severity of the impairment, rather than the chronicity, could determine the effects of the MT in the motor impairment.

### Sensory function

Although no significant improvement was detected in any group regarding kinesthesia and stereognosis, the MT provided a statistically significant improvement in the tactile sensation in comparison to the results obtained from participants in the experimental group,

mainly promoted by an improvement in the ability to sense light touches. Despite the chronicity and the limited duration of the treatment, the effect size of the difference between groups for the tactile sensation subscale was moderate. Furthermore, a detailed analysis of its items showed a considerable effect size for light touch sensitivity. The authors hypothesize that these improvements can be related to the visual enhancement of touch, which suggests that tactile perception could be augmented by viewing the stimulated parts.<sup>31</sup> Visual feedback can exert a strong modulatory influence over the motor system when there is a conflict, which can, in turn, override other modalities, such as proprioception.<sup>32</sup> Neuroimaging techniques have shown the dominance of vision over proprioception during motor programming.<sup>32</sup> Interestingly, this mechanism has previously been confirmed in healthy and brain damaged individuals with lower somatosensory sensitivity,<sup>31</sup> as is the case in the present study. The participants were required to pay special attention to the intervention task, which could have been another factor that contributed to the improvement in tactile sensitivity. Attention to touch could have led to increased activation in somatosensory cortical areas, including the primary somatosensory cortex.<sup>33</sup> Interestingly, the role of attention in MT has been previously reported to have a positive impact on hemineglect.<sup>10, 16, 34</sup>

An improvement in temperature sensation was also detected in our study but without showing differences between groups, and a similar result had been previously reported in a stroke population with mild to moderate paresis after MT.<sup>14</sup> This change may have been attributed to multimodal neurons in the posterior parietal and premotor cortical areas that respond to sensory stimuli, which modulate the somatosensory cortex network and contribute to the recovery of the somatosensory system. The severity of the impairment could have influenced the different results.

### Limitations

The limitations of this study must be taken into account when analyzing the results. First, the sample size, which consisted of 31 participants, can be considered small even though it is similar to other studies.<sup>10, 14</sup> Second, the chronicity and the severity of the motor impairment restricts the extrapolation of the results to other

population, because these factors seem to determine the effects of the MT, as described throughout the text. Third, the progress of the participants was not determined in a follow-up assessment. Finally, the MCID of the NSA scale has not been validated, which complicates the evaluation of the functional impact of the significant changes detected in this scale.

Nonetheless, the improvement in light touch sensitivity after the MT intervention may be relevant, because somatosensory deficits occur in approximately 50% of stroke survivors and can limit functional recovery.<sup>31</sup> MT could therefore be a tool to enhance sensory function in very chronic individuals with severely affected upper limb function.

### Conclusions

Our results suggest that MT can provide limited but positive effects in light touch sensitivity in chronic stroke survivors with severely impaired upper limb function while providing similar motor improvement. Our results and previous research in the field suggest that MT may be more effective for motor improvement in mild to moderate hemiparesis, even at chronic stages, rather than in severe paresis, where MT may provide benefits to tactile sensitivity.

### References

1. Invernizzi M, Negrini S, Carda S, Lanzotti L, Cisari C, Baricich A. The value of adding mirror therapy for upper limb motor recovery of subacute stroke patients: a randomized controlled trial. *Eur J Phys Rehabil Med* 2013;49:311-7.
2. Park Y, Chang M, Kim KM, An DH. The effects of mirror therapy with tasks on upper extremity function and self-care in stroke patients. *J Phys Ther Sci*. May 2015;27:1499-501.
3. Kwakkel G, Kollen BJ, van der Grond J, Prevo AJ. Probability of regaining dexterity in the flaccid upper limb: impact of severity of paresis and time since onset in acute stroke. *Stroke* 2003;34:2181-6.
4. Barker RN, Gill TJ, Brauer SG. Factors contributing to upper limb recovery after stroke: a survey of stroke survivors in Queensland Australia. *Disabil Rehabil* 2007;29:981-9.
5. Nestor ABM, Jamie Bitensky M, Katherine Salter BA, Robert Teasel Md F. The Role of Task-Specific Training in Rehabilitation Therapies. *Top Stroke Rehabil* 2005;12:58-65.
6. Lum PS, Mulroy S, Amdur RL, Requejo P, Prilutsky BI, Dromerick AW. Gains in upper extremity function after stroke via recovery or compensation: Potential differential effects on amount of real-world limb use. *Top Stroke Rehabil* 2009;16:237-53.
7. Lindberg PG, Schmitz C, Engardt M, Forssberg H, Borg J. Use-dependent up- and down-regulation of sensorimotor brain circuits in stroke patients. *Neurorehabil Neural Repair* 2007;21:315-26.
8. Taub E, Uswatte G, Mark VW, Morris DM. The learned nonuse phenomenon: implications for rehabilitation. *Eura Medicophys* 2006;42:241-56.

9. Deconinck FJ, Smorenburg AR, Benham A, Ledebt A, Feltham MG, Savelsbergh GJ. Reflections on Mirror Therapy: A Systematic Review of the Effect of Mirror Visual Feedback on the Brain. *Neurorehabil Neural Repair* 2015;29:349-61.
10. Dohle C, Pullen J, Nakaten A, Kust J, Rietz C, Karbe H. Mirror therapy promotes recovery from severe hemiparesis: a randomized controlled trial. *Neurorehabil Neural Repair* 2009;23:209-17.
11. Michielsen ME, Smits M, Ribbers GM, et al. The neuronal correlates of mirror therapy: an fMRI study on mirror induced visual illusions in patients with stroke. *J Neurol Neurosurg Psychiatry* 2011;82:393-8.
12. Michielsen ME, Selles RW, van der Geest JN, et al. Motor recovery and cortical reorganization after mirror therapy in chronic stroke patients: a phase II randomized controlled trial. *Neurorehabil Neural Repair* 2011;25:223-33.
13. Rossiter HE, Borrelli MR, Borchert RJ, Bradbury D, Ward NS. Cortical Mechanisms of Mirror Therapy After Stroke. *Neurorehabil Neural Repair* 2015;29:444-52.
14. Wu CY, Huang PC, Chen YT, Lin KC, Yang HW. Effects of mirror therapy on motor and sensory recovery in chronic stroke: a randomized controlled trial. *Arch Phys Med Rehabil* 2013;94:1023-30.
15. Park JY, Chang M, Kim KM, Kim HJ. The effect of mirror therapy on upper-extremity function and activities of daily living in stroke patients. *J Phys Ther Sci* 2015;27:1681-3.
16. Thieme H, Bayn M, Wurg M, Zange C, Pohl M, Behrens J. Mirror therapy for patients with severe arm paresis after stroke--a randomized controlled trial. *Clin Rehabil* 2013;27:314-24.
17. Shah SK. Reliability of the Original Brunnstrom Recovery Scale Following Hemiplegia. *Aust Occup Ther J* 1984;31:144-51.
18. Sanford J, Moreland J, Swanson LR, Stratford PW, Gowland C. Reliability of the Fugl-Meyer assessment for testing motor performance in patients following stroke. *Phys Ther* 1993;73:447-54.
19. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189-98.
20. Romero M, Sanchez A, Marin C, Navarro MD, Ferri J, Noe E. [Clinical usefulness of the Spanish version of the Mississippi Aphasia Screening Test (MASTsp): validation in stroke patients]. *Neurologia* 2012;27:216-24 [Article in Spanish].
21. Lincoln NB, Jackson JM, Adams SA. Reliability and Revision of the Nottingham Sensory Assessment for Stroke Patients. *Physiotherapy* 1998;84:358-65.
22. Lang CE, Edwards DF, Birkenmeier RL, Dromerick AW. Estimating minimal clinically important differences of upper-extremity measures early after stroke. *Arch Phys Med Rehabil* 2008;89:1693-700.
23. Fregni F, Pascual-Leone A. Hand motor recovery after stroke: tuning the orchestra to improve hand motor function. *Cogn Behav Neurol* 2006;19:21-33.
24. Kwakkel G, Kollen BJ, van der Grond J, Prevo AJH. Probability of Regaining Dexterity in the Flaccid Upper Limb: Impact of Severity of Paresis and Time Since Onset in Acute Stroke. *Stroke* 2003;34:2181-6.
25. Kong K-H, Chua KSG, Lee J. Recovery of upper limb dexterity in patients more than 1 year after stroke: Frequency, clinical correlates and predictors. *NeuroRehabilitation* 2011;28:105-11.
26. Coupar F, Pollock A, Rowe P, Weir C, Langhorne P. Predictors of upper limb recovery after stroke: a systematic review and meta-analysis. *Clin Rehabil* 2012;26:291-313.
27. Lee YY, Hsieh YW, Wu CY, Lin KC, Chen CK. Proximal Fugl-Meyer Assessment Scores Predict Clinically Important Upper Limb Improvement After Three Stroke Rehabilitative Interventions. *Arch Phys Med Rehabil* 2015;96:2137-44.
28. Patel AT, Duncan PW, Lai S-M, Studenski S. The relation between impairments and functional outcomes poststroke. *Arch Phys Med Rehabil* 2000;81:1357-63.
29. Mercier C, Bourbonnais D. Relative shoulder flexor and handgrip strength is related to upper limb function after stroke. *Clin Rehabil* 2004;18:215-21.
30. Samuelkamaleshkumar S, Reethajanetsureka S, Pauljebaraj P, Ben-shamir B, Padankatti SM, David JA. Mirror therapy enhances motor performance in the paretic upper limb after stroke: a pilot randomized controlled trial. *Arch Phys Med Rehabil* 2014;95:2000-5.
31. Serino A, Farnè A, Rinaldesi ML, Haggard P, Ládavas E. Can vision of the body ameliorate impaired somatosensory function? *Neuropsychologia* 2007;45:1101-7.
32. Touzalin-Chretien P, Ehrler S, Dufour A. Dominance of vision over proprioception on motor programming: evidence from ERP. *Cereb Cortex* 2010;20:2007-16.
33. Johansen-Berg H, Christensen V, Woolrich M, Matthews PM. Attention to touch modulates activity in both primary and secondary somatosensory areas. *NeuroReport* 2000;11:1237-41.
34. Pandian JD, Arora R, Kaur P, Sharma D, Vishwambaran DK, Arima H. Mirror therapy in unilateral neglect after stroke (MUST trial): a randomized controlled trial. *Neurology* 2014;83:1012-7.

*Funding.*—The study presented in the manuscript was conducted under the PhD Program in Medicine of the Universitat Autònoma de Barcelona and was funded in part by Ministerio de Economía y Competitividad of Spain (Project NeuroVR, TIN2013-44741-R and Project REACT, TIN2014-61975-EXP) and by Universitat Politècnica de València (Grant PAID-10-14).

*Conflicts of interest.*—The authors certify that there is no conflict of interest with any financial organization regarding the material discussed in the manuscript. Article first published online: February 29, 2016. - Manuscript accepted: February 2, 2016. - Manuscript revised: January 15, 2016. - Manuscript received: October 15, 2015.